

Vegetation Patterns in Semi-Arid Environments

Jonathan A. Sherratt

Department of Mathematics
Heriot-Watt University

CMPD3, Bordeaux, May/June 2010

This talk can be downloaded from my web site

`www.ma.hw.ac.uk/~jas`

Outline

- 1 Ecological Background
- 2 The Mathematical Model
- 3 Linear Analysis
- 4 Travelling Wave Equations
- 5 Conclusions

Vegetation Pattern Formation



Bushy vegetation in Niger



Mitchell grass in Australia

(Western New South Wales)


- Banded vegetation patterns are found on gentle slopes in semi-arid areas of Africa, Australia and Mexico
- First identified by aerial photos in 1950s
- Plants vary from grasses to shrubs and trees

Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water


Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes



Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes



- The stripes move uphill (very slowly)

Two Key Ecological Questions

- How does the spacing of the vegetation bands depend on rainfall, herbivory and slope?
- At what rainfall level is there a transition to desert?

Outline

- 1 Ecological Background
- 2 The Mathematical Model
- 3 Linear Analysis
- 4 Travelling Wave Equations
- 5 Conclusions

Mathematical Model of Klausmeier

Rate of change = Rainfall – Evaporation – Uptake by + Flow
of water plants downhill

Rate of change = Growth, proportional – Mortality + Random
plant biomass to water uptake dispersal

$$\partial w / \partial t = A - w - wu^2 + \nu \partial w / \partial x$$

$$\partial u / \partial t = wu^2 - Bu + \partial^2 u / \partial x^2$$

Mathematical Model of Klausmeier

Rate of change = Rainfall – Evaporation – Uptake by + Flow
of water plants downhill

Rate of change = Growth, proportional – Mortality + Random
plant biomass to water uptake dispersal

$$\partial w / \partial t = A - w - wu^2 + \nu \partial w / \partial x$$

$$\partial u / \partial t = wu^2 - Bu + \partial^2 u / \partial x^2$$

The nonlinearity in wu^2 arises because the presence of roots increases water infiltration into the soil.

Mathematical Model of Klausmeier

Rate of change = Rainfall – Evaporation – Uptake by + Flow
of water plants downhill


Rate of change = Growth, proportional – Mortality + Random
plant biomass to water uptake dispersal

$$\partial w / \partial t = A - w - wu^2 + \nu \partial w / \partial x$$


$$\partial u / \partial t = wu^2 - Bu + \partial^2 u / \partial x^2$$

Parameters: A : rainfall B : plant loss ν : slope


Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model



Typical Solution of the Model



Typical Solution of the Model



Outline

- 1 Ecological Background
- 2 The Mathematical Model
- 3 Linear Analysis**
- 4 Travelling Wave Equations
- 5 Conclusions

Homogeneous Steady States

- For all parameter values, there is a stable “desert” steady state $u = 0$, $w = A$

Homogeneous Steady States


- For all parameter values, there is a stable “desert” steady state $u = 0, w = A$
- When $A \geq 2B$, there are also two non-trivial steady states, one of which is unstable to homogeneous perturbations

Homogeneous Steady States

- For all parameter values, there is a stable “desert” steady state $u = 0, w = A$
- When $A \geq 2B$, there are also two non-trivial steady states, one of which is unstable to homogeneous perturbations
- Patterns develop when the other steady state (u_s, w_s) is unstable to inhomogeneous perturbations

Approximate Conditions for Patterning


Look for solutions $(u, w) = (u_s, w_s) + (u_0, w_0) \exp\{ikx + \lambda t\}$



The dispersion relation $\text{Re}[\lambda(k)]$ is algebraically complicated

Approximate Conditions for Patterning

Look for solutions $(u, w) = (u_s, w_s) + (u_0, w_0) \exp\{ikx + \lambda t\}$




The dispersion relation $\text{Re}[\lambda(k)]$ is algebraically complicated

An approximate condition for pattern formation is

$$A < \nu^{1/2} B^{5/4} / 8^{1/4}$$

Approximate Conditions for Patterning

Look for solutions $(u, w) = (u_s, w_s) + (u_0, w_0) \exp\{ikx + \lambda t\}$




The dispersion relation $\text{Re}[\lambda(k)]$ is algebraically complicated

An approximate condition for pattern formation is

$$2B < A < \nu^{1/2} B^{5/4} / 8^{1/4}$$


One can naively assume that existence of (u_s, w_s) gives a second condition

An Illustration of Conditions for Patterning



The dots show parameters for which there are growing linear modes.


An Illustration of Conditions for Patterning



Numerical simulations show patterns in both the dotted and green regions of parameter space.


Predicting Pattern Wavelength

Pattern wavelength is the most accessible property of vegetation stripes in the field, via aerial photography. Wavelength can be predicted from the linear analysis



Predicting Pattern Wavelength

Pattern wavelength is the most accessible property of vegetation stripes in the field, via aerial photography. Wavelength can be predicted from the linear analysis



However this prediction doesn't fit the patterns seen in numerical simulations.

Shortcomings of Linear Stability Analysis

Linear stability analysis fails in two ways:

- It significantly over-estimates the minimum rainfall level for patterns.
- Close to the maximum rainfall level for patterns, it incorrectly predicts a variation in pattern wavelength with rainfall.

Outline

- 1 Ecological Background
- 2 The Mathematical Model
- 3 Linear Analysis
- 4 Travelling Wave Equations**
- 5 Conclusions

Travelling Wave Equations

The patterns move at constant shape and speed


$$\Rightarrow u(x, t) = U(z), w(x, t) = W(z), z = x - ct$$

$$d^2U/dz^2 + c dU/dz + WU^2 - BU = 0$$


$$(\nu + c)dW/dz + A - W - WU^2 = 0$$

The patterns are periodic (limit cycle) solutions of these equations

When do Patterns Form?





Pattern Formation for Low Rainfall





Patterns are also seen for parameters in the green region.


Pattern Formation for Low Rainfall




Pattern Formation for Low Rainfall



Minimum Rainfall for Patterns




Pattern Stability



The wavelengths shown are those compatible with periodic boundary conditions on a domain of length 80.

Pattern Stability



The wavelengths shown are those compatible with periodic boundary conditions on a domain of length 80.


Pattern Stability

Key Result

Many of the possible patterns are unstable and thus will never be seen.



However, for a wide range of rainfall levels, there are multiple stable patterns.

Hysteresis





- The existence of multiple stable patterns raises the possibility of hysteresis
- We consider slow variations in the rainfall parameter A
- Parameters correspond to grass, and the rainfall range corresponds to 130–930 mm/year

Hysteresis




<< Mode 5 >> <<<< Mode 1 >>>> < Mode 3 >

Hysteresis



<< Mode 5 >> <<<< Mode 1 >>>>> < Mode 3 >

Wavelength vs Rainfall




Outline

- 1 Ecological Background
- 2 The Mathematical Model
- 3 Linear Analysis
- 4 Travelling Wave Equations
- 5 Conclusions

Predictions of Pattern Wavelength


- In general, pattern wavelength depends on initial conditions
- When vegetation stripes arise from homogeneous vegetation via a decrease in rainfall, pattern wavelength will remain at its bifurcating value.



Predictions of Pattern Wavelength

- In general, pattern wavelength depends on initial conditions
- When vegetation stripes arise from homogeneous vegetation via a decrease in rainfall, pattern wavelength will remain at its bifurcating value.

$$\text{Wavelength} = \sqrt{\frac{8\pi^2}{B_V}}$$



References

J.A. Sherratt: An analysis of vegetation stripe formation in semi-arid landscapes. *J. Math. Biol.* 51, 183-197 (2005).

J.A. Sherratt, G.J. Lord: Nonlinear dynamics and pattern bifurcations in a model for vegetation stripes in semi-arid environments. *Theor. Pop. Biol.* 71, 1-11 (2007).



J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments I. Submitted.

J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments II. Patterns with the largest possible propagation speeds. Submitted.

List of Frames

1

Ecological Background

- Vegetation Pattern Formation
- Mechanisms for Vegetation Patterning
- Two Key Ecological Questions

2

The Mathematical Model

- Mathematical Model of Klausmeier
- Typical Solution of the Model

3

Linear Analysis

- Homogeneous Steady States
- Approximate Conditions for Patterning
- An Illustration of Conditions for Patterning
- Predicting Pattern Wavelength
- Shortcomings of Linear Stability Analysis

4

Travelling Wave Equations

- Travelling Wave Equations
- When do Patterns Form?
- Pattern Formation for Low Rainfall
- Pattern Stability
- Hysteresis

5

Conclusions

- Predictions of Pattern Wavelength
- References