

Nonlinear Dynamics of Vegetation Patterns in Semi-Deserts

Jonathan A. Sherratt

Department of Mathematics
Heriot-Watt University

University of Dundee
26 May 2011

This talk can be downloaded from my web site

www.ma.hw.ac.uk/~jas

Outline

- 1 Ecological Background
- 2 The Mathematical Model
- 3 Travelling Wave Equations
- 4 Pattern Stability
- 5 Conclusions

Vegetation Pattern Formation



Bushy vegetation in Niger



Mitchell grass in Australia

(Western New South Wales)


- Banded vegetation patterns are found on gentle slopes in semi-arid areas of Africa, Australia and Mexico
- First identified by aerial photos in 1950s
- Plants vary from grasses to shrubs and trees

Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water


Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes




Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes



Mechanisms for Vegetation Patterning

- Basic mechanism: competition for water
- Possible detailed mechanism: water flow downhill causes stripes



- The stripes move uphill (very slowly)

Two Key Ecological Questions

- How does the spacing of the vegetation bands depend on rainfall, herbivory and slope?
- At what rainfall level is there a transition to desert?

Outline

- 1 Ecological Background
- 2 The Mathematical Model**
- 3 Travelling Wave Equations
- 4 Pattern Stability
- 5 Conclusions

Mathematical Model of Klausmeier

Rate of change = Rainfall – Evaporation – Uptake by + Flow
of water plants downhill

Rate of change = Growth, proportional – Mortality + Random
plant biomass to water uptake dispersal

$$\partial w / \partial t = A - w - wu^2 + v \partial w / \partial x$$

$$\partial u / \partial t = wu^2 - Bu + \partial^2 u / \partial x^2$$

Mathematical Model of Klausmeier

Rate of change = Rainfall – Evaporation – Uptake by + Flow
of water plants downhill

Rate of change = Growth, proportional – Mortality + Random
plant biomass to water uptake dispersal

$$\partial w / \partial t = A - w - wu^2 + v \partial w / \partial x$$

$$\partial u / \partial t = wu^2 - Bu + \partial^2 u / \partial x^2$$

The nonlinearity in wu^2 arises because the presence of roots increases water infiltration into the soil.

Mathematical Model of Klausmeier

Rate of change = Rainfall – Evaporation – Uptake by + Flow
of water plants downhill


Rate of change = Growth, proportional – Mortality + Random
plant biomass to water uptake dispersal

$$\partial w / \partial t = A - w - wu^2 + \nu \partial w / \partial x$$


$$\partial u / \partial t = wu^2 - Bu + \partial^2 u / \partial x^2$$

Parameters: **A:** rainfall **B:** plant loss **ν :** slope


Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model



Typical Solution of the Model



Typical Solution of the Model



Homogeneous Steady States

- For all parameter values, there is a stable “desert” steady state $u = 0$, $w = A$

Homogeneous Steady States


- For all parameter values, there is a stable “desert” steady state $u = 0, w = A$
- When $A \geq 2B$, there are also two non-trivial steady states, one of which is unstable to homogeneous perturbations

Homogeneous Steady States

- For all parameter values, there is a stable “desert” steady state $u = 0, w = A$
- When $A \geq 2B$, there are also two non-trivial steady states, one of which is unstable to homogeneous perturbations
- Patterns develop when the other steady state (u_s, w_s) is unstable to inhomogeneous perturbations

Approximate Conditions for Patterning


Look for solutions $(u, w) = (u_s, w_s) + (u_0, w_0) \exp\{ikx + \lambda t\}$



The dispersion relation $\text{Re}[\lambda(k)]$ is algebraically complicated

Approximate Conditions for Patterning

Look for solutions $(u, w) = (u_s, w_s) + (u_0, w_0) \exp\{ikx + \lambda t\}$




The dispersion relation $\text{Re}[\lambda(k)]$ is algebraically complicated

An approximate condition for pattern formation is

$$A < \nu^{1/2} B^{5/4} / 8^{1/4}$$

Approximate Conditions for Patterning

Look for solutions $(u, w) = (u_s, w_s) + (u_0, w_0) \exp\{ikx + \lambda t\}$




The dispersion relation $\text{Re}[\lambda(k)]$ is algebraically complicated

An approximate condition for pattern formation is

$$2B < A < \nu^{1/2} B^{5/4} / 8^{1/4}$$


One can naively assume that existence of (u_s, w_s) gives a second condition

An Illustration of Conditions for Patterning



The dots show parameters for which there are growing linear modes.


An Illustration of Conditions for Patterning



Numerical simulations show patterns in both the dotted and green regions of parameter space.


Predicting Pattern Wavelength

Pattern wavelength is the most accessible property of vegetation stripes in the field, via aerial photography. Wavelength can be predicted from the linear analysis.



Predicting Pattern Wavelength

Pattern wavelength is the most accessible property of vegetation stripes in the field, via aerial photography. Wavelength can be predicted from the linear analysis.



However this prediction doesn't fit the patterns seen in numerical simulations.

Shortcomings of Linear Stability Analysis

Linear stability analysis fails in two ways:

- It significantly over-estimates the minimum rainfall level for patterns.
- Close to the maximum rainfall level for patterns, it incorrectly predicts a variation in pattern wavelength with rainfall.

Outline

- 1 Ecological Background
- 2 The Mathematical Model
- 3 Travelling Wave Equations**
- 4 Pattern Stability
- 5 Conclusions

Travelling Wave Equations

The patterns move at constant shape and speed


$$\Rightarrow u(x, t) = U(z), w(x, t) = W(z), z = x - ct$$

$$d^2U/dz^2 + c dU/dz + WU^2 - BU = 0$$



$$(\nu + c)dW/dz + A - W - WU^2 = 0$$

The patterns are periodic (limit cycle) solutions of these equations


Bifurcation Diagram for Travelling Wave Equations




Bifurcation Diagram for Travelling Wave Equations



When do Patterns Form?





Pattern Formation for Low Rainfall





Patterns are also seen for parameters in the green region.


Pattern Formation for Low Rainfall



Pattern Formation for Low Rainfall



Minimum Rainfall for Patterns



Outline

- 1 Ecological Background
- 2 The Mathematical Model
- 3 Travelling Wave Equations
- 4 Pattern Stability**
- 5 Conclusions

The Eigenvalue Problem

$$\text{PDE model: } \begin{aligned} u_t &= u_{zz} + cu_z + f(u, w) \\ w_t &= \nu w_z + cw_z + g(u, w) \end{aligned}$$

$$\begin{aligned} \text{Periodic wave satisfies: } 0 &= U_{zz} + cU_z + f(U, W) \\ 0 &= \nu W_z + cW_z + g(U, W) \end{aligned}$$

$$\begin{aligned} \text{Consider } u(z, t) &= U(z) + e^{\lambda t} \bar{u}(z) \quad \text{with } |\bar{u}| \ll |U| \\ w(z, t) &= W(z) + e^{\lambda t} \bar{w}(z) \quad \text{with } |\bar{w}| \ll |W| \end{aligned}$$

$$\begin{aligned} \Rightarrow \text{Eigenfunction eqn: } \lambda \bar{u} &= \bar{u}_{zz} + c\bar{u}_z + f_u(U, W)\bar{u} + f_w(U, W)\bar{w} \\ \lambda \bar{w} &= \nu \bar{w}_z + c\bar{w}_z + g_u(U, W)\bar{u} + g_w(U, W)\bar{w} \end{aligned}$$

$$\begin{aligned} \text{Boundary conditions: } \bar{u}(0) &= \bar{u}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \\ \bar{w}(0) &= \bar{w}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \end{aligned}$$

The Eigenvalue Problem

$$\begin{aligned} \text{PDE model: } u_t &= u_{zz} + cu_z + f(u, w) \\ w_t &= \nu w_z + cw_z + g(u, w) \end{aligned}$$

Periodic wave satisfies:

$$\begin{aligned} 0 &= U_{zz} + cU_z + f(U, W) \\ 0 &= \nu W_z + cW_z + g(U, W) \end{aligned}$$

Consider

$$\begin{aligned} u(z, t) &= U(z) + e^{\lambda t} \bar{u}(z) \quad \text{with } |\bar{u}| \ll |U| \\ w(z, t) &= W(z) + e^{\lambda t} \bar{w}(z) \quad \text{with } |\bar{w}| \ll |W| \end{aligned}$$

$$\begin{aligned} \Rightarrow \text{Eigenfunction eqn: } \lambda \bar{u} &= \bar{u}_{zz} + c\bar{u}_z + f_u(U, W)\bar{u} + f_w(U, W)\bar{w} \\ \lambda \bar{w} &= \nu \bar{w}_z + c\bar{w}_z + g_u(U, W)\bar{u} + g_w(U, W)\bar{w} \end{aligned}$$

Boundary conditions:

$$\begin{aligned} \bar{u}(0) &= \bar{u}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \\ \bar{w}(0) &= \bar{w}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \end{aligned}$$

The Eigenvalue Problem

$$\begin{aligned} \text{PDE model: } u_t &= u_{zz} + cu_z + f(u, w) \\ w_t &= \nu w_z + cw_z + g(u, w) \end{aligned}$$

$$\begin{aligned} \text{Periodic wave satisfies: } 0 &= U_{zz} + cU_z + f(U, W) \\ 0 &= \nu W_z + cW_z + g(U, W) \end{aligned}$$

Consider $u(z, t) = U(z) + e^{\lambda t} \bar{u}(z)$ with $|\bar{u}| \ll |U|$
 $w(z, t) = W(z) + e^{\lambda t} \bar{w}(z)$ with $|\bar{w}| \ll |W|$

$$\begin{aligned} \Rightarrow \text{Eigenfunction eqn: } \lambda \bar{u} &= \bar{u}_{zz} + c\bar{u}_z + f_u(U, W)\bar{u} + f_w(U, W)\bar{w} \\ \lambda \bar{w} &= \nu \bar{w}_z + c\bar{w}_z + g_u(U, W)\bar{u} + g_w(U, W)\bar{w} \end{aligned}$$

$$\begin{aligned} \text{Boundary conditions: } \bar{u}(0) &= \bar{u}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \\ \bar{w}(0) &= \bar{w}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \end{aligned}$$

The Eigenvalue Problem

$$\begin{aligned} \text{PDE model: } u_t &= u_{zz} + cu_z + f(u, w) \\ w_t &= \nu w_z + cw_z + g(u, w) \end{aligned}$$

$$\begin{aligned} \text{Periodic wave satisfies: } 0 &= U_{zz} + cU_z + f(U, W) \\ 0 &= \nu W_z + cW_z + g(U, W) \end{aligned}$$

$$\begin{aligned} \text{Consider } u(z, t) &= U(z) + e^{\lambda t} \bar{u}(z) \quad \text{with } |\bar{u}| \ll |U| \\ w(z, t) &= W(z) + e^{\lambda t} \bar{w}(z) \quad \text{with } |\bar{w}| \ll |W| \end{aligned}$$

$$\begin{aligned} \Rightarrow \text{Eigenfunction eqn: } \lambda \bar{u} &= \bar{u}_{zz} + c\bar{u}_z + f_u(U, W)\bar{u} + f_w(U, W)\bar{w} \\ \lambda \bar{w} &= \nu \bar{w}_z + c\bar{w}_z + g_u(U, W)\bar{u} + g_w(U, W)\bar{w} \end{aligned}$$

$$\begin{aligned} \text{Boundary conditions: } \bar{u}(0) &= \bar{u}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \\ \bar{w}(0) &= \bar{w}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \end{aligned}$$

The Eigenvalue Problem

$$\begin{aligned} \text{PDE model: } u_t &= u_{zz} + cu_z + f(u, w) \\ w_t &= \nu w_z + cw_z + g(u, w) \end{aligned}$$

$$\begin{aligned} \text{Periodic wave satisfies: } 0 &= U_{zz} + cU_z + f(U, W) \\ 0 &= \nu W_z + cW_z + g(U, W) \end{aligned}$$

$$\begin{aligned} \text{Consider } u(z, t) &= U(z) + e^{\lambda t} \bar{u}(z) \quad \text{with } |\bar{u}| \ll |U| \\ w(z, t) &= W(z) + e^{\lambda t} \bar{w}(z) \quad \text{with } |\bar{w}| \ll |W| \end{aligned}$$

$$\begin{aligned} \Rightarrow \text{Eigenfunction eqn: } \lambda \bar{u} &= \bar{u}_{zz} + c\bar{u}_z + f_u(U, W)\bar{u} + f_w(U, W)\bar{w} \\ \lambda \bar{w} &= \nu \bar{w}_z + c\bar{w}_z + g_u(U, W)\bar{u} + g_w(U, W)\bar{w} \end{aligned}$$

$$\begin{aligned} \text{Boundary conditions: } \bar{u}(0) &= \bar{u}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \\ \bar{w}(0) &= \bar{w}(L)e^{i\gamma} \quad (0 \leq \gamma < 2\pi) \end{aligned}$$

The Eigenvalue Problem


$$\begin{aligned}\text{Eigenfunction eqn: } \lambda \bar{u} &= \bar{u}_{zz} + c\bar{u}_z + f_u(U, W)\bar{u} + f_w(U, W)\bar{w} \\ \lambda \bar{w} &= \nu\bar{w}_z + c\bar{w}_z + g_u(U, W)\bar{u} + g_w(U, W)\bar{w}\end{aligned}$$

Here $0 < z < L$, with $(\bar{u}, \bar{w})(0) = (\bar{u}, \bar{w})(L)e^{i\gamma}$ ($0 \leq \gamma < 2\pi$)


The Eigenvalue Problem

$$\begin{aligned} \text{Eigenfunction eqn: } \lambda \bar{u} &= \bar{u}_{zz} + c\bar{u}_z + f_u(U, W)\bar{u} + f_w(U, W)\bar{w} \\ \lambda \bar{w} &= \nu\bar{w}_z + c\bar{w}_z + g_u(U, W)\bar{u} + g_w(U, W)\bar{w} \end{aligned}$$

Here $0 < z < L$, with $(\bar{u}, \bar{w})(0) = (\bar{u}, \bar{w})(L)e^{i\gamma}$ ($0 \leq \gamma < 2\pi$)




$$\text{Re}(\lambda) < 0$$




The Eigenvalue Problem

$$\begin{aligned} \text{Eigenfunction eqn: } \lambda \bar{u} &= \bar{u}_{zz} + c\bar{u}_z + f_u(U, W)\bar{u} + f_w(U, W)\bar{w} \\ \lambda \bar{w} &= \nu\bar{w}_z + c\bar{w}_z + g_u(U, W)\bar{u} + g_w(U, W)\bar{w} \end{aligned}$$

Here $0 < z < L$, with $(\bar{u}, \bar{w})(0) = (\bar{u}, \bar{w})(L)e^{i\gamma}$ ($0 \leq \gamma < 2\pi$)



$$\text{Re}(\lambda) > 0$$



Numerical Calculation of Eigenvalue Spectrum

(based on Jens Rademacher, Bjorn Sandstede, Arnd Scheel Physica D 229 166-183, 2007)


- 1 solve numerically for the periodic wave by continuation in c from a Hopf bifurcation point in the travelling wave eqns

$$\begin{aligned}0 &= U_{zz} + cU_z + f(U, W) \\0 &= \nu W_z + cW_z + g(U, W) \quad (z = x - ct)\end{aligned}$$

Numerical Calculation of Eigenvalue Spectrum

(based on Jens Rademacher, Bjorn Sandstede, Arnd Scheel Physica D 229 166-183, 2007)

- 1 solve numerically for the periodic wave by continuation in c from a Hopf bifurcation point in the travelling wave equations
- 2 for $\gamma = 0$, discretise the eigenfunction equations in space, giving a (large) matrix eigenvalue problem




$$\lambda \bar{u} = \bar{u}_{zz} + c \bar{u}_z + f_u(U, W) \bar{u} + f_w(U, W) \bar{w}, \quad \bar{u}(0) = \bar{u}(L) e^{i\gamma}$$

$$\lambda \bar{w} = \nu \bar{w}_z + c \bar{w}_z + g_u(U, W) \bar{u} + g_w(U, W) \bar{w}, \quad \bar{w}(0) = \bar{w}(L) e^{i\gamma}$$

Numerical Calculation of Eigenvalue Spectrum

(based on Jens Rademacher, Bjorn Sandstede, Arnd Scheel Physica D 229 166-183, 2007)

- 1 solve numerically for the periodic wave by continuation in c from a Hopf bifurcation point in the travelling wave equations
- 2 for $\gamma = 0$, discretise the eigenfunction equations in space, giving a (large) matrix eigenvalue problem
- 3 continue the eigenfunction equations numerically in γ , starting from each of the periodic eigenvalues




$$\begin{aligned} \lambda \bar{u} &= \bar{u}_{zz} + c \bar{u}_z + f_u(U, W) \bar{u} + f_w(U, W) \bar{w}, & \bar{u}(0) &= \bar{u}(L) e^{i\gamma} \\ \lambda \bar{w} &= \nu \bar{w}_z + c \bar{w}_z + g_u(U, W) \bar{u} + g_w(U, W) \bar{w}, & \bar{w}(0) &= \bar{w}(L) e^{i\gamma} \end{aligned}$$

Numerical Calculation of Eigenvalue Spectrum

(based on Jens Rademacher, Bjorn Sandstede, Arnd Scheel Physica D 229 166-183, 2007)


- 1 solve numerically for the periodic wave by continuation in c from a Hopf bifurcation point in the travelling wave equations
- 2 for $\gamma = 0$, discretise the eigenfunction equations in space, giving a (large) matrix eigenvalue problem
- 3 continue the eigenfunction equations numerically in γ , starting from each of the periodic eigenvalues



This gives the eigenvalue spectrum, and hence (in)stability


Numerical Calculation of Eigenvalue Spectrum

(based on Jens Rademacher, Bjorn Sandstede, Arnd Scheel Physica D 229 166-183, 2007)



STABLE

Eckhaus
instability



UNSTABLE

This gives the eigenvalue spectrum, and hence (in)stability

Stability in a Parameter Plane


By following this procedure at each point on a grid in parameter space, regions of stability/instability can be determined.

In fact, stable/unstable boundaries can be computed accurately by numerical continuation of the point at which

$$\operatorname{Re}\lambda = \operatorname{Im}\lambda = \gamma = \partial^2 \operatorname{Re}\lambda / \partial \gamma^2 = 0$$

(Eckhaus instability point)

Stability in a Parameter Plane




Pattern Stability: The Key Result

Key Result

Many of the possible patterns are unstable and thus will never be seen.


However, for a wide range of rainfall levels, there are multiple stable patterns.

Hysteresis




- The existence of multiple stable patterns raises the possibility of hysteresis
- We consider slow variations in the rainfall parameter A
- Parameters correspond to grass, and the rainfall range corresponds to 130–930 mm/year

Hysteresis




Time

Space




<< Mode 5 >> <<<<< Mode 1 >>>>> < Mode 3 >

Hysteresis




Time

Space



<< Mode 5 >> <<<<< Mode 1 >>>>> < Mode 3 >

Wavelength vs Rainfall




Outline

- 1 Ecological Background
- 2 The Mathematical Model
- 3 Travelling Wave Equations
- 4 Pattern Stability
- 5 Conclusions**

Predictions of Pattern Wavelength


- In general, pattern wavelength depends on initial conditions
- When vegetation stripes arise from homogeneous vegetation via a decrease in rainfall, pattern wavelength will remain at its bifurcating value.



Predictions of Pattern Wavelength

- In general, pattern wavelength depends on initial conditions
- When vegetation stripes arise from homogeneous vegetation via a decrease in rainfall, pattern wavelength will remain at its bifurcating value.

$$\text{Wavelength} = \sqrt{\frac{8\pi^2}{B\nu}}$$



References

- J.A. Sherratt: An analysis of vegetation stripe formation in semi-arid landscapes. *J. Math. Biol.* 51, 183-197 (2005).
- J.A. Sherratt, G.J. Lord: Nonlinear dynamics and pattern bifurcations in a model for vegetation stripes in semi-arid environments. *Theor. Pop. Biol.* 71, 1-11 (2007).
- J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments I. *Nonlinearity* 23, 2657-2675 (2010).
- J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments II. Patterns with the largest possible propagation speeds. Submitted.
- J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments III. The transition between homoclinic solutions. Submitted.

List of Frames

- 1 Ecological Background**
 - Vegetation Pattern Formation
 - Mechanisms for Vegetation Patterning
 - Two Key Ecological Questions
- 2 The Mathematical Model**
 - Mathematical Model of Klausmeier
 - Typical Solution of the Model
 - Homogeneous Steady States
 - Approximate Conditions for Patterning
 - Shortcomings of Linear Stability Analysis
- 3 Travelling Wave Equations**
 - Travelling Wave Equations
 - Bifurcation Diagram for Travelling Wave Equations
 - When do Patterns Form?
 - Pattern Formation for Low Rainfall
- 4 Pattern Stability**
 - The Eigenvalue Problem
 - Numerical Calculation of Eigenvalue Spectrum
 - Stability in a Parameter Plane
 - Hysteresis

- 5 Conclusions**
 - Predictions of Pattern Wavelength
 - References