

History-Dependent Patterns of Whole Ecosystems

Jonathan A. Sherratt


Department of Mathematics
and Maxwell Institute for Mathematical Sciences
Heriot-Watt University

OCCAM, Oxford, 1-5 July 2013


This talk can be downloaded from my web site

`www.ma.hw.ac.uk/~jas`

Vegetation Patterns



1950



(William MacFadyden,
Geogr. J. 115: 199-211, 1950)

- 1 Ecological Background
- 2 A Simple Mathematical Model
- 3 Travelling Wave Equations
- 4 Pattern Stability
- 5 Other Examples of Landscape-Scale Patterns

Vegetation Patterns



Bushy vegetation in Niger




Mitchell grass in Australia

(Western New South Wales)

- Banded vegetation patterns are found on gentle slopes in semi-arid areas of Africa, Australia and Mexico
- Plants vary from grasses to shrubs and trees
- Typical wavelength 1km for shrubs and trees

Why Do Plants Form Patterns?



Data from Burkina Faso

Rietkerk et al

Plant Ecology 148: 207-224, 2000


More plants \Rightarrow more roots and organic matter in soil
 \Rightarrow more infiltration of rainwater

Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill


Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill




Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill



Banded Patterns on Slopes

- On slopes, water flow downhill causes stripes which move uphill



Key Ecological Questions

- At what rainfall level is there a switch from uniform vegetation to patterns?
- At what rainfall level is there a transition to desert?
- How does the spacing of the vegetation bands depend on rainfall, herbivory and slope?

Outline

- 1 Ecological Background
- 2 **A Simple Mathematical Model**
- 3 Travelling Wave Equations
- 4 Pattern Stability
- 5 Other Examples of Landscape-Scale Patterns

Mathematical Model of Klausmeier

Rate of change = Rainfall – Evaporation – Uptake by + Flow
of water plants downhill

Rate of change = Growth, proportional – Mortality + Random
plant biomass to water uptake dispersal

$$\partial w / \partial t = A - w - wu^2 + \nu \partial w / \partial x$$

$$\partial u / \partial t = wu^2 - Bu + \partial^2 u / \partial x^2$$

Mathematical Model of Klausmeier

Rate of change of water = Rainfall – Evaporation – Uptake by plants + Flow downhill


Rate of change of plant biomass = Growth, proportional to water uptake – Mortality + Random dispersal

$$\partial w / \partial t = A - w - wu^2 + \nu \partial w / \partial x$$

$$\partial u / \partial t = wu^2 - Bu + \partial^2 u / \partial x^2$$

The nonlinearity in wu^2 arises because the presence of plants increases water infiltration into the soil.

Mathematical Model of Klausmeier



$$wu^2 = w \cdot u \cdot (\text{infiltration rate})$$

The nonlinearity in wu^2 arises because the presence of plants increases water infiltration into the soil.

Mathematical Model of Klausmeier

Rate of change = Rainfall – Evaporation – Uptake by + Flow
of water plants downhill


Rate of change = Growth, proportional – Mortality + Random
plant biomass to water uptake dispersal

$$\partial w / \partial t = A - w - wu^2 + \nu \partial w / \partial x$$


$$\partial u / \partial t = wu^2 - Bu + \partial^2 u / \partial x^2$$

Parameters: A : rainfall B : plant loss ν : slope


Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model




Typical Solution of the Model



Typical Solution of the Model



Typical Solution of the Model



Homogeneous Steady States

- For all parameter values, there is a stable “desert” steady state $u = 0$, $w = A$

Homogeneous Steady States


- For all parameter values, there is a stable “desert” steady state $u = 0, w = A$
- When $A \geq 2B$, there are also two non-trivial steady states, one of which is unstable to homogeneous perturbations

Homogeneous Steady States

- For all parameter values, there is a stable “desert” steady state $u = 0, w = A$
- When $A \geq 2B$, there are also two non-trivial steady states, one of which is unstable to homogeneous perturbations
- The other steady state (u_s, w_s) is stable to homogeneous perturbations but can be unstable to inhomogeneous perturbations \Rightarrow pattern formation

Approximate Conditions for Patterning


Look for solutions $(u, w) = (u_s, w_s) + (u_0, w_0) \exp\{ikx + \lambda t\}$



The dispersion relation $\text{Re}[\lambda(k)]$ is algebraically complicated

Approximate Conditions for Patterning

Look for solutions $(u, w) = (u_s, w_s) + (u_0, w_0) \exp\{ikx + \lambda t\}$



The dispersion relation $\text{Re}[\lambda(k)]$ is algebraically complicated

To leading order for large ν , the condition for pattern formation is

$$A < B^{5/4} \nu^{1/2} (\sqrt{2} - 1)^{1/2}$$

Back to Key Ecological Questions

- At what rainfall level is there a switch from uniform vegetation to patterns?
- At what rainfall level is there a transition to desert?
- How does the spacing of the vegetation bands depend on rainfall, herbivory and slope?

Outline

- 1 Ecological Background
- 2 A Simple Mathematical Model
- 3 Travelling Wave Equations**
- 4 Pattern Stability
- 5 Other Examples of Landscape-Scale Patterns

Travelling Wave Equations

The patterns move at constant shape and speed


$$\Rightarrow u(x, t) = U(z), w(x, t) = W(z), z = x - ct$$

$$d^2 U/dz^2 + c dU/dz + WU^2 - BU = 0$$



$$(\nu + c)dW/dz + A - W - WU^2 = 0$$

The patterns are periodic (limit cycle) solutions of these equations


Bifurcation Diagram for Travelling Wave Equations



Bifurcation Diagram for Travelling Wave Equations




When do Patterns Form?



Pattern Formation for Low Rainfall

Min rainfall
for patterns

Turing
bifurcation




Min rainfall
for uniform veg


Pattern Formation for Low Rainfall

Min rainfall
for patterns

Turing
bifurcation



Min rainfall
for uniform veg



Back to Key Ecological Questions


- At what rainfall level is there a switch from uniform vegetation to patterns?
- At what rainfall level is there a transition to desert?
- How does the spacing of the vegetation bands depend on rainfall, herbivory and slope?

Outline

- 1 Ecological Background
- 2 A Simple Mathematical Model
- 3 Travelling Wave Equations
- 4 Pattern Stability**
- 5 Other Examples of Landscape-Scale Patterns


Pattern Stability

Not all of the possible patterns are stable as solutions of the model equations.




Pattern Stability: Numerical Approach

The boundary between stable and unstable patterns can be calculated by numerical continuation of the essential spectrum.




Calculations of this type can be performed using the software package WAVETRAN (www.ma.hw.ac.uk/wavetrain).

Pattern Stability: Wavelength vs Rainfall



The wavelengths shown are those compatible with periodic boundary conditions on a domain of length 80.

Pattern Stability: Wavelength vs Rainfall



The wavelengths shown are those compatible with periodic boundary conditions on a domain of length 80.

Pattern Stability: The Key Result



Key Result

Many of the possible patterns are unstable and thus will never be seen.

However, for a wide range of rainfall levels, there are multiple stable patterns.



Variations in Rainfall: Hysteresis

The existence of multiple stable patterns suggests the possibility of hysteresis.



Domain length 150, periodic bc's

Data on the Effects of Changing Rainfall



Data from 1950-1995 (C. Valentin & J.M. d'Herbès, Catena 37:231, 1999)


Back to Key Ecological Questions

- At what rainfall level is there a switch from uniform vegetation to patterns?
- At what rainfall level is there a transition to desert?
- How does the spacing of the vegetation bands depend on rainfall, herbivory and slope?

Predictions of Pattern Wavelength

- In general, pattern wavelength depends on initial conditions
- When vegetation stripes arise from homogeneous vegetation via a decrease in rainfall, pattern wavelength will remain at its bifurcating value.


$$\text{Wavelength} = \sqrt{\frac{8\pi^2}{B_V}}$$



Outline


- 1 Ecological Background
- 2 A Simple Mathematical Model
- 3 Travelling Wave Equations
- 4 Pattern Stability
- 5 Other Examples of Landscape-Scale Patterns

Tree Patches in Savannah Grasslands



(Olivier Lejeune et al, Phys. Rev. E 66: 010901, 2002)

Pattern of Fog-Dependent Vegetation in Chile



Tillandsia landbeckii

Aerial photo over Atacama Desert, Northern Chile
(Borthagaray et al, J. Theor. Biol. 265: 18-26, 2010)

Ribbon Forest in Colorado, USA



Photo taken by David Buckner

Mudflat Pattern in The Netherlands



(Weerman et al, Am. Nat. 176: E15-E32, 2010)

Mussel Bed Pattern in the Wadden Sea


In the Wadden Sea, mussel beds self-organise into striped patterns



Aerial photo of
a mussel bed

Mussel Bed Pattern in the Wadden Sea

In the Wadden Sea, mussel beds self-organise into striped patterns



Aerial photo of
a mussel bed

References

- J.A. Sherratt: An analysis of vegetation stripe formation in semi-arid landscapes. *J. Math. Biol.* 51, 183-197 (2005).
- J.A. Sherratt, G.J. Lord: Nonlinear dynamics and pattern bifurcations in a model for vegetation stripes in semi-arid environments. *Theor. Pop. Biol.* 71, 1-11 (2007).
- J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments I. *Nonlinearity* 23, 2657-2675 (2010).
- J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments II: patterns with the largest possible propagation speeds. *Proc. R. Soc. Lond. A* 467, 3272-3294 (2011).
- J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments III: the transition between homoclinic solutions. *Physica D* 242, 30-41 (2013).
- J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments IV: slowly moving patterns and their stability. *SIAM J. Appl. Math.* 73, 330-350 (2013).
- J.A. Sherratt: History-dependent patterns of whole ecosystems. *Ecological Complexity* 14, 8-20 (2013).
- J.A. Sherratt: Pattern solutions of the Klausmeier model for banded vegetation in semi-arid environments V: the transition from patterns to desert. *SIAM J. Appl. Math.* in press.

List of Frames

- 1 **Ecological Background**
 - Vegetation Patterns
 - Why Do Plants Form Patterns?
 - Banded Patterns on Slopes
 - Key Ecological Questions
- 2 **A Simple Mathematical Model**
 - Mathematical Model of Klausmeier
 - Typical Solution of the Model
 - Homogeneous Steady States
 - Approximate Conditions for Patterning
 - Back to Key Ecological Questions
- 3 **Travelling Wave Equations**
 - Travelling Wave Equations
 - Bifurcation Diagram for Travelling Wave Equations
 - When do Patterns Form?
 - Pattern Formation for Low Rainfall

- 4 **Pattern Stability**
 - Pattern Stability
 - Variations in Rainfall: Hysteresis
 - Predictions of Pattern Wavelength
- 5 **Other Examples of Landscape-Scale Patterns**
 - Photo Gallery of Landscape-Scale Patterns
 - References

Mussel Bed Pattern in the Wadden Sea


In the Wadden Sea, mussel beds self-organise into striped patterns



Aerial photo of
a mussel bed

Mussel Bed Pattern in the Wadden Sea

In the Wadden Sea, mussel beds self-organise into striped patterns



Aerial photo of
a mussel bed

Mussel Bed Pattern in the Wadden Sea

Model of van de Koppel *et al*

(Am Nat 165: E66, 2005)

$a(x, t)$ = density of algae

$m(x, t)$ = density of mussels

$$\begin{aligned}
 \frac{\partial a}{\partial t} &= \overbrace{\beta \frac{\partial a}{\partial x}}^{\text{advection by tide}} + \overbrace{\alpha(1-a)}^{\text{transfer to/from upper water layers}} - \overbrace{am}^{\text{eaten by mussels}} \\
 \frac{\partial m}{\partial t} &= \underbrace{\frac{\partial^2 m}{\partial x^2}}_{\text{random movement}} + \underbrace{\delta am}_{\text{birth}} - \underbrace{\gamma \frac{m}{(1+m)}}_{\text{dislodgement by waves}}
 \end{aligned}$$



Aerial photo of
 a mussel bed

Mussel Bed Pattern in the Wadden Sea

Model of van de Koppel *et al*

(Am Nat 165: E66, 2005)

$a(x, t)$ = density of algae


$m(x, t)$ = density of mussels

$$\begin{aligned}
 \frac{\partial a}{\partial t} &= \underbrace{\beta \frac{\partial a}{\partial x}}_{\text{advection by tide}} + \underbrace{\alpha(1-a)}_{\text{transfer to/from upper water layers}} - \underbrace{am}_{\text{eaten by mussels}} \\
 \frac{\partial m}{\partial t} &= \underbrace{\frac{\partial^2 m}{\partial x^2}}_{\text{random movement}} + \underbrace{\delta am}_{\text{birth}} - \underbrace{\gamma \frac{m}{(1+m)}}_{\text{dislodgement by waves}}
 \end{aligned}$$




Aerial photo of
 a mussel bed

Typical Pattern Solution



Pattern Existence and Stability



The parameter δ reflects the supply rate of algae

Hysteresis in Mussel Bed Patterns

