

Civil Engineering 2 Mathematics Autumn 2011

Exercise Sheet 5

1. Verify the relation

$$\nabla \times (\nabla \times \mathbf{A}) = \nabla(\nabla \cdot \mathbf{A}) - \tilde{\nabla}^2 \mathbf{A}$$

for the vector field

$$\mathbf{A} = x^2 y \mathbf{i} - 2xz \mathbf{j} + 2yz \mathbf{k}.$$

(Use the definition $\tilde{\nabla}^2 \mathbf{A} = \partial^2 \mathbf{A} / \partial x^2 + \partial^2 \mathbf{A} / \partial y^2 + \partial^2 \mathbf{A} / \partial z^2$).

2. If $\mathbf{A} = 2yz \mathbf{i} - x^2 y \mathbf{j} + xz^2 \mathbf{k}$, $\mathbf{B} = x^2 \mathbf{i} + yz \mathbf{j} - xy \mathbf{k}$ and $\phi = 2x^2 yz^3$, find:

(i) $(\mathbf{A} \cdot \nabla)\phi$; (ii) $\mathbf{A} \cdot (\nabla\phi)$; (iii) $(\mathbf{B} \cdot \nabla)\mathbf{A}$; (iv) $\mathbf{A} \times (\nabla\phi)$; (v) $(\mathbf{A} \times \nabla)\phi$.

(Note that in (iii) there is some abuse of notation.) Verify that the answers to (i) & (ii) and (iv) & (v) are the same.

3. Calculate the divergence and curl of the following vector fields:

(i) $\mathbf{v} = \mathbf{r}$, where $\mathbf{r} = x \mathbf{i} + y \mathbf{j} + z \mathbf{k}$;

(ii) $\mathbf{v} = \frac{\mathbf{r}}{|\mathbf{r}|}$ with \mathbf{r} as in (i);

(iii) $\mathbf{v} = \omega \mathbf{k} \times \mathbf{r}$, where $\mathbf{r} = x \mathbf{i} + y \mathbf{j}$ and ω is a constant;

(iv) $\mathbf{v} = \omega \mathbf{k} \times \frac{\mathbf{r}}{|\mathbf{r}|}$, with \mathbf{r} and ω as in (iii).

4. Establish the identity

$$\nabla \cdot (\phi \mathbf{G}) = \phi (\nabla \cdot \mathbf{G}) + (\nabla \phi) \cdot \mathbf{G},$$

for a general scalar field $\phi(x, y, z)$ and general vector field $\mathbf{G} = G_1 \mathbf{i} + G_2 \mathbf{j} + G_3 \mathbf{k}$. Hence evaluate

$$\nabla \cdot \left(\frac{y}{xr^3} \mathbf{r} \right),$$

where $\mathbf{r} = (x, y, z)$ and $r = |\mathbf{r}|$.

5. If $\mathbf{A} = A_1 \mathbf{i} + A_2 \mathbf{j} + A_3 \mathbf{k}$ is a vector and ϕ is a scalar, prove that

(i) $\nabla \times (\nabla \phi) = 0$;

(ii) $\nabla \cdot (\nabla \times \mathbf{A}) = 0$.

Answers:

1. Both right hand side and left hand side are equal to $(2x + 2)\mathbf{j}$.

2. (i) $8xy^2z^4 - 2x^4yz^3 + 6x^3yz^4$; (ii) same as (i); (iii) $(2yz^2 - 2xy^2)\mathbf{i} - (x^2yz + 2x^3y)\mathbf{j} + (x^2z^2 - 2x^2yz)\mathbf{k}$;

(iv) $-(6x^4y^2z^2 + 2x^3z^5)\mathbf{i} + (4x^2yz^5 - 12x^2y^2z^3)\mathbf{j} + (4x^2yz^4 + 4x^3y^2z^3)\mathbf{k}$;
(v) same as (iv).

3. (i) $\nabla \cdot \mathbf{v} = 3, \nabla \times \mathbf{v} = 0$; (ii) $\nabla \cdot \mathbf{v} = 2/r, \nabla \times \mathbf{v} = 0$; (iii) $\nabla \cdot \mathbf{v} = 0, \nabla \times \mathbf{v} = 2\omega \mathbf{k}$;

(iv) $\nabla \cdot v = 0, \nabla \times \mathbf{v} = (\omega/r)k$.

4. $\nabla \cdot \left(\frac{y}{xr^3}\mathbf{r}\right) = 0$.